Effects of a community intervention to reduce the serving of alcohol to intoxicated patrons

Author(s)

Warpenius K, Holmila M, Mustonen H

Published

2010

Publisher

Addiction

Type

Journal article

Volume

105

Issue

6

Page(s)

1032-40

Abstract

Aims To assess the effects of an alcohol prevention programme to reduce the serving of alcoholic beverages to intoxicated clients on licensed premises. Research design A controlled pre- (2004) and post-intervention study (2006) design. Intervention A community-based programme combining law enforcement, responsible beverage service training, information campaigns and policy initiatives in one Finnish town (Jyvaskyla). Participants and measurements A male actor pretended to be clearly under the influence of alcohol and tried to buy a pint of beer at licensed premises. For the baseline measurement, every bar and nightclub was visited in the intervention and the control areas (94 licensed premises in total). Post-intervention data were gathered with the same principles (100 licensed premises in total). A researcher observed every visit and documented the results. Results In the post-intervention study there was a statistically significant increase in refusals to serve denials alcohol to the actor in the intervention area (from 23% to 42% of the licensed premises) compared to refusals in the control area (from 36% to 27% of the licensed premises). Conclusion Previous research has documented thatmulti-component communitybased interventions can have a significant impact on over-serving of alcohol when training and house policies are combined with effective law inforcement. The present findings also demonstrate that comprehensive Responsible Beverage Service (RBS) interventions applied at a local community level can be effective in decreasing service to intoxicated clients in a Nordic context.

Web Link

 $\underline{http://www.ncbi.nlm.nih.gov/pubmed/20219058}$

View PDF