

The detection and prevention of unintentional consumption of DOx and 25x-NBOMe at Portugal's Boom Festival

Author(s)

Martins D, Barratt M, Vale Pires C, Carvalho H, Ventura M, Fornís I, Valente H

Published

2017

Publisher

Human Psychopharmacology: Clinical and experimental

Type

Journal article

Volume

32

Issue

3

Abstract

Objective:

This paper describes the misrepresentation of LSD at Portugal's Boom Festival 2014 and the prevention of unintentional consumption of DOx and 25x-NBOMe among LSD consumers attending a drug-checking service.

Methods:

Two hundred forty-five drug samples expected to contain LSD were submitted to the drug-checking service for chemical analysis. One hundred ten post-test questionnaires were successfully matched with test results.

Results:

About 67.3% of the alleged LSD samples tested contained only LSD; 0.8% contained LSD combined with adulterants; 24.1% did not contain LSD but did contain another psychoactive substance, including 11.4% that were 2,5-dimethoxyamphetamine derivatives and 9.8% that were N-benzyl-2,5-dimethoxyphenethylamine derivatives; and no psychoactive substance was detected in 7.8%. The majority of service users who received unexpected test results regarding their alleged LSD (74.2%) reported that they did not intend to consume the drug. Following dissemination of alerts on day 2, a larger than expected proportion of all tests conducted were for LSD, when comparing the 2014 festival to 2012, where no such alert was disseminated.

Conclusions:

Although these results support the provision of integrated drug-checking services in party settings, evidence of their utility and effectiveness would be improved through future research incorporating more robust measures of outcomes following provision of drug-checking results.

Web Link

[Link to the article](#)

[View PDF](#)