Polysubstance use and associated effects at raves parties

Author(s)

Fermín Fernández-Calderón; Óscar M. Lozano Rojas; Izaskun Bilbao Acedos; Antonio J. Rojas Tejada; Claudio Vidal Giné; Esperanza Vergara Moragues & Francisco González-Saiz

Published

2012

Publisher

Health and Addictions

Type

Journal article

Volume

12

Issue

1

Page(s)

37-59

Abstract

Objetives:

The purpose of this study is to describe positive and negative effects perceived by underground rave attenders, and to analyze the relations between drugs consume and perceived risks.

Method:

The research was developed with a sample of 252 assistant at 22 underground raves in Andalucía (Spain), from May to October 2008. It has been applied contrast proportions and logistic regressions analysis.

Results:

Results show that the most consumed drugs were: alcohol (94.4%), cannabis

(76.6%), speed (74.7%), ecstasy (64.1%), cocaine (28.2%), ketamine (23.8%) and LSD (22.9%). The most positive effects perceived were: euphoria, well-being, happiness (90.3%) and connection with music (75.4%). The most negative effects perceived were: tiredness (78.2%), jaw clenching (55.2%) and concentration decreased (54.4%). Ecstasy is the substance which more increases probability to perceive positive effects, while speed is the one that more increases probability to perceive negative effects.

Conclusions:

Negative effects experimented for participants show the high risk for rave attendees' health. These, together with positive effects, must be considered in order to design preventive and risk reduction strategies.

Web Link
Link to the article
View PDF